Extremal Sets Minimizing Dimension–normalized Boundary in Hamming Graphs

M. Cemil Azizoğlu* and Ömer Eğecioğlu†
Department of Computer Science
University of California at Santa Barbara
{azizoglu, omer}@cs.ucsb.edu

Abstract

We prove that the set of first k vertices of a Hamming graph in reverse–lexicographic order constitutes an extremal set minimizing the dimension–normalized edge–boundary over all k–vertex subsets of the graph. This generalizes a result of Lindsey and can be used to prove a tight lower bound for the isoperimetric number and the bisection width of arrays.

Keywords: Hamming graph, product graph, array, isoperimetric number, bisection width, extremal set, partition, Schur-convexity, majorization.

1 Introduction

In this paper, we consider questions of the following general form: Given a graph G and a natural number k, what is the optimum value of a certain quantity in a set of k vertices of G? The desired quantity could be the number of edges between a set of k vertices and its complement (i.e. the size of the boundary), or the number of edges induced by a set of k vertices, etc. The sets achieving the optimum value are called *extremal sets*.

Specifically, we study extremal sets in Hamming graphs minimizing the size of the edge–boundary of a set of vertices of given size, where boundary edges along each dimension are *normalized* by a weight determined by that dimension as shall soon be explained.

First, we introduce some notation and terminology. Given a graph G and a subset X of its vertices, let ∂X denote the *edge–boundary*, or simply *boundary*, of X. This is the set of edges connecting vertices in X with vertices not in X (i.e. the complement of X). A *d–dimensional Hamming graph* H^d is a graph with $k_1 \times k_2 \times \cdots \times k_d$ vertices, $k_1 \leq k_2 \leq \cdots \leq k_d$, each having a unique label $l = \langle l_1, l_2, \cdots, l_d \rangle$ where $0 \leq l_i \leq k_i - 1$. There is an edge between two vertices iff their labels differ in exactly one digit. A *d–dimensional array* A^d resembles H^d with the exception that two vertices are adjacent if and only if their labels differ in exactly one digit *and* the difference is

*Supported in part by NSF Grant No. CCR–9821038 and by a fellowship from The İzmir Institute of Technology, İzmir, Turkey.

†Supported in part by NSF Grant No. CCR–9821038.

exactly one. Examples of a 2–dimensional Hamming graph and a 2–dimensional array are shown in Figure 1.



Figure 1: The 2–dimensional Hamming graph $K_3 \times K_4$ and array $P_3 \times P_4$.

The *Cartesian product* $G \times H$ of two graphs G and H is the graph with vertex set $V(G) \times V(H)$, in which vertices (u, v) and (u', v') are adjacent iff u is adjacent to u' in G and $v = v'$, or v is adjacent to v' in H and $u = u'$. The constituent graphs G and H are called *factors*. A Hamming graph can be characterized as the Cartesian product of a number of *complete graphs* of different sizes, i.e. $H^d = K_{k_1} \times K_{k_2} \times \cdots \times K_{k_d}$ where K_r is a complete graph on r vertices. Similarly, A^d can be characterized as the Cartesian product of a number of *path graphs* of varying length, i.e. $A^d = P_{k_1} \times P_{k_2} \times \cdots \times P_{k_d}$ where P_r is a path graph (chain) with r vertices.

Lindsey [19] proved that the set of first k vertices of a Hamming graph in *lexicographic order* constitutes an extremal set minimizing the boundary ∂X over all k–element subsets X. The lexicographic order is defined as follows: In the Hamming graph $H^d = K_{k_1} \times K_{k_2} \times \cdots \times K_{k_d}$ with $k_1 \leq k_2 \leq \cdots \leq k_d$, vertex $x = \langle x_1, \cdots, x_d \rangle$ precedes vertex $y = \langle y_1, \cdots, y_d \rangle$ in lexicographic order iff there exists an index i such that $x_1 = y_1$, $x_2 = y_2$, ..., $x_{i-1} = y_{i-1}$ and $x_i < y_i$ holds. Intuitively, in lexicographic order, we traverse the Hamming graph in the direction of the *next largest factor* starting with the vertex labelled $\langle 0, 0, \cdots, 0 \rangle$. For instance, the vertices of the Hamming graph in Figure 1 in lexicographic order are labelled

$$00,\ 01,\ 02,\ 03,\ 10,\ 11,\ 12,\ 13,\ 20,\ 21,\ 22,\ 23.$$

Our aim in this paper is to determine and describe extremal sets of Hamming graphs minimizing the dimension–normalized boundary. This is defined next.

Definition 1 *Given a Hamming graph $H^d = K_{k_1} \times K_{k_2} \times \cdots \times K_{k_d}$ and a subset X of its vertices,* the dimension–normalized boundary $B(X)$ *of X is defined as*

$$B(X) = \frac{|\partial_1 X|}{c_1} + \frac{|\partial_2 X|}{c_2} + \cdots + \frac{|\partial_d X|}{c_d} \tag{1}$$

where for $1 \leq i \leq d$, $\partial_i X$ *is the set of boundary edges along dimension* i *and*

$$c_i = \begin{cases} k_i^2 & if\ k_i\ is\ even, \\ k_i^2 - 1 & if\ k_i\ is\ odd. \end{cases} \tag{2}$$

We prove that the set of first k vertices in *reverse–lexicographic order* constitutes an extremal set minimizing the dimension–normalized boundary over all k–element subsets in a Hamming graph. The definition of the reverse–lexicographic order is similar to that of the lexicographic order: In the Hamming graph $H^d = K_{k_1} \times K_{k_2} \times \cdots \times K_{k_d}$ with $k_1 \leq k_2 \leq \cdots \leq k_d$, vertex $x = \langle x_1, \cdots, x_d \rangle$ precedes vertex $y = \langle y_1, \cdots, y_d \rangle$ in reverse–lexicographic order iff there exists an index i such that $x_d = y_d$, $x_{d-1} = y_{d-1}$, ..., $x_{i+1} = y_{i+1}$ and $x_i < y_i$ holds. In other words, we move in the direction of the *next smallest factor* starting at the vertex labelled $\langle 0, 0, \cdots, 0 \rangle$. To illustrate, the vertices of the Hamming graph in the above example listed in reverse–lexicographic order are

$$00,\ 10,\ 20,\ 01,\ 11,\ 21,\ 02,\ 12,\ 22,\ 03,\ 13,\ 23.$$

We should point out that there are other sets of vertices which are structurally equivalent to the sets specified in our definitions of lexicographic or reverse–lexicographic orders. These are obtained by symmetries in the underlying graph. For instance, in Figure 1, another ordering structurally equivalent to the lexicographic ordering would be 23, 22, 21, 20, 13 etc.. Similarly, the sets defined by the initial segments of the ordering 23, 13, 03, 22, 12, etc., give rise to sets structurally identical to those in reverse–lexicographic order.

We state our claim formally in the following theorem.

Theorem 1 *Given a* d*–dimensional Hamming graph* H^d*, let* X *be any* k*–vertex subset of* $V(H^d)$ *and* $\overline{X}$ *be the set of first* k *vertices of* H^d *in reverse–lexicographic order. Then* $B(\overline{X}) \leq B(X)$.

Interestingly, when all factors of H^d have equal size, the lexicographic and reverse–lexicographic orders both result in structurally symmetric subsets and hence are equivalent with respect to extremal sets minimizing the boundary (dimension–normalized or otherwise). Therefore Theorem 1 is trivially true when $k_1 = k_2 = \cdots = k_d$ by Lindsey's result, since the denominators c_i in (1) will all be equal and minimizing $B(X)$ will be equivalent to minimizing $|\partial_1 X| + |\partial_2 X| + \cdots + |\partial_d X| = |\partial X|$, i.e. the size of the boundary of X.

In the next section, we describe the notion of the isoperimetric number, which is a quantity closely related to extremal sets. The isoperimetric number problem for special classes of graphs provides the basis of our motivation for this work.

1.1 Motivation

An important quantity in the theory of graphs is the *isoperimetric number* $i(G)$ of a graph G, defined as

$$i(G) = \min_{1 \leq |X| \leq \frac{|V(G)|}{2}} \frac{|\partial X|}{|X|} \tag{3}$$

where $X \subseteq V(G)$. That is, the set of vertices of G is partitioned into two nonempty sets and the ratio of the number of edges between the two parts and the number of vertices in the smaller one is minimized. A subset X achieving the equality in (3) is called an *isoperimetric set*.

The notion of the isoperimetric number of a graph G serves as a measure of connectivity of G as it quantifies the minimal interaction between a set of vertices X and its complement $V(G) \setminus X$ in terms of the number of edges between them. In many instances, the isoperimetric number of a graph can be used to obtain a tight lower bound for its *bisection width* as well. We refer the reader to Mohar [22] or Chung [12] for a discussion of basic results and various interesting properties of $i(G)$.

At present, the isoperimetric number of an array $A^d = P_{k_1} \times P_{k_2} \times \cdots \times P_{k_d}$ is known only when either $k_1 = k_2 = \cdots = k_d$ (see Azizoğlu and Eğecioğlu [4]), or the size of the largest factor is even (see Azizoğlu and Eğecioğlu [5]. The latter is also implicit in [10] ([17])). The techniques used to obtain these results seem to fail in the general case. However using the notion of extremal sets minimizing dimension–normalized boundary, one can show that

$$i(A^d) = \min_i \frac{1}{\lfloor \frac{k_i}{2} \rfloor}$$

The technique used involves embedding a Hamming graph into A^d and associating these extremal sets with isoperimetric sets of the array. We refer the reader to [6] for details.

1.2 A summary of previous results

There has been a significant amount of research in the area of isoperimetric bounds on various popular classes of graphs such as Hamming graphs, arrays and tori. We shall only mention those results in this area which pertain to our discussion and refer the reader to Bezrukov [8] for a comprehensive survey and Bollobás [9] for a general discussion of this and related topics.

As mentioned before, an extremal set of a graph for a given k is, in a broad sense, a configuration of k vertices with

- minimum number of boundary edges, or
- maximum number of spanned edges

among all such k–vertex subsets of the given graph. The problem of finding extremal sets of the first (or, second) type is called *the minimum–boundary–edge problem* (or, *the maximum–induced–edge problem*). It can be shown that the minimum–boundary–edge and the maximum–induced–edge problems are equivalent for regular graphs [11]. We remark that one can easily obtain the isoperimetric number of a given graph if the extremal sets of the first type are known (and the boundary is actually computable). Evidently, an extremal set X with $\lfloor |V(G)|/2 \rfloor$ vertices in a given graph G determines a bisection for G.

The maximum–induced–edge problem (hence the minimum–boundary–edge problem, because of its regularity) for the hypercube (d–dimensional binary Hamming graph) was solved by Harper [14] and extended by Lindsey [19] to the d–dimensional k–ary Hamming graph. In both instances, there is a nested structure of solutions, and the first k vertices in *lexicographic order* constitute an extremal set. The maximum–induced–edge problem for the d–dimensional k–ary array A_k^d was solved by Bollobás and Leader [11]. Since A_k^d is not regular, this result does not automatically give a solution to the minimum–boundary–edge problem. It was later extended to general arrays by Ahlswede and Bezrukov [1] who also gave a solution for $P_{k_1} \times P_{k_2}$ for the minimum–boundary–edge problem. The first nontrivial bounds on the minimum–boundary–edge problem for the

d–dimensional k–ary arrays are in Bollobás and Leader [11]. Unfortunately, however, the bounds obtained are not tight enough to yield an exact formula for $i(A_k^d)$.

Similar problems have been studied in the literature for the vertex–boundary of a given configuration of vertices. For instance, for the d–dimensional k–ary torus, Bollobás and Leader [10] solved the vertex–boundary problem for even k. Riordan [24] later extended their result by giving an ordering of vertices, on the d–dimensional even torus, which minimizes the number of vertices at shortest distance t from the vertices in the ordering. Wang and Wang [25] solved this problem for $P_\infty \times \cdots \times P_\infty$, i.e. the *$d$–dimensional infinite array*, where the minimum is taken over all nonempty finite subsets of vertices. In their result, each P_∞ may be infinite in both directions or in one direction only. They also gave a simple ordering of the vertices in which the first k vertices constitute an extremal set minimizing the vertex–boundary. In a recent paper, Harper [15] solved the vertex–boundary problem on Hamming graphs.

1.3 Outline

The outline of the remainder of this paper is as follows. In Section 2 we consider the case of two–dimensional Hamming graphs. First we define the terminology we use and state a number of basic facts on restricted integer partitions, majorization, and Schur-convexity. Then we identify potential extremal sets in H^2 as integer partitions inside a rectangle. The problem of showing that the set of first k vertices of H^2 in reverse–lexicographic order constitutes an extremal set minimizing the dimension–normalized edge–boundary over all k–vertex subsets, becomes the problem of maximization of a certain function on partitions, which is a linear combination of two Schur-convex functions. However the function itself is not Schur-convex, and the identification of the partition on which the maximum is achieved is actually done using an inductive argument. The main result of this section is Lemma 1. In Section 3 we extend the proof to the higher–dimensional case. This is done by an induction on the number of dimensions, using the two–dimensional result as the base case. Finally, concluding remarks are given in Section 4.

2 The two–dimensional case

Let $H^2 = K_m \times K_n$ be a given 2–dimensional Hamming graph. Without loss of generality, we may assume that $m = k_1 \leq k_2 = n$. Consider a subset X of vertices in H^2. Let X' be the subset of vertices of H^2 obtained by pushing (compressing) all the vertices in X as far downward, and then to the left in H^2 as possible. It is easy to see (and proved in [19], [16]) that $B(X') \leq B(X)$ since the number of boundary edges in either dimension will not increase as a result of this procedure. A subset X' in the compressed form corresponds to a *partition* of the integer $|X|$ contained in the $m \times n$ rectangle.

We give below the definitions and properties of partitions that we will use in our proof of Theorem 1. The reader is referred to [2] for further details.

Partitions

A *partition* λ of an integer N is a sequence $(\lambda_1, \lambda_2, \ldots, \lambda_\ell)$ of positive integers (called *parts*) satisfying $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_\ell$ and $\lambda_1 + \lambda_2 + \cdots + \lambda_\ell = N$. We put $|\lambda| = N$. The Ferrers diagram of λ is a 2–dimensional array of unit cells (or nodes) in which row i from the bottom has λ_i cells,

and the rows are left justified. It is clear that in our case, $\lambda = X'$ forms a partition of $|X|$ whose Ferrers diagram is contained in the $m \times n$ rectangle, i.e. $\lambda_i \leq m$ for $1 \leq i \leq \ell$ (i.e. each part at most m), and $\ell \leq n$ (i.e. number of parts at most n). We use $\mathbb{P}(m,n)$ to denote the set of these partitions. Thus we may assume that an extremal set is a partition $\lambda \in \mathbb{P}(m,n)$, and use the symbol $\mathbb{P}(m,n)$ to refer to $H^2 = K_m \times K_n$ when we are not interested in the graph structure of H^2, but just the placement of the subset λ. We may augment partitions by adding parts of zero length and write $\sum_{i \geq 1} \lambda_i$ for $|\lambda|$. We also identify partitions with their diagrams when there is no confusion.



Figure 2: The partition $\lambda = (5,4,3,3,1,1)$.

Given a partition $\lambda = (\lambda_1, \lambda_2, \ldots, \lambda_n)$, we may define a new partition $\lambda' = (\lambda'_1, \lambda'_2, \ldots, \lambda'_m)$ by choosing λ'_i as the number of parts of λ that are $\geq i$. The partition λ' is called the *conjugate* of λ. Geometrically, λ' is obtained from λ by reflection in the main diagonal (equivalently by counting the cells in successive columns of λ). For example, the conjugate of $(5,4,3,3,1,1)$ is $(6,4,4,2,1)$. Clearly $|\lambda| = |\lambda'|$, and if $\lambda \in \mathbb{P}(m,n)$ then $\lambda' \in \mathbb{P}(n,m)$.

Majorization, Schur-convexity, and transfer

As is to be remarked later in the proof, the concept of majorization and Schur-convexity are not needed to prove the claim. However, their relevance shall be pointed out here anyway, due to their importance.

Given two partitions $\lambda = (\lambda_1, \lambda_2, \ldots, \lambda_N)$ and $\mu = (\mu_1, \mu_2, \ldots, \mu_N)$ of N, λ is *majorized* by μ, written $\lambda \prec \mu$, if

$$\lambda_1 + \lambda_2 + \cdots + \lambda_k \leq \mu_1 + \mu_2 + \cdots + \mu_k, \qquad k = 1, 2, \ldots, N.$$

Majorization is also referred to as the *dominance* or *natural* order ([20], Ch. 1). As soon as $N \geq 6$, majorization is not a total ordering. For example the partitions $(3,1,1,1)$, and $(2,2,2)$ of 6 are not comparable. However reverse–lexicographic ordering on partitions is a linear extension of $\prec$. Thus

$$\lambda \prec \mu \;\Rightarrow\; \lambda \leq \mu.$$

Furthermore $\lambda \prec \mu \;\Leftrightarrow\; \mu' \prec \lambda'$ (see [20], (1.11)). A real valued function g defined on partitions of an integer N is said to be *Schur–convex* (see [21], Ch. 3) if

$$\lambda \prec \mu \;\Rightarrow\; g(\lambda) \leq g(\mu).$$

We make use of the following special case of a result of Schur, 1923; Hardy, Littlewood, and Polya, 1929 (see [21], Ch 3, Proposition C.1)

Proposition 1 *Suppose ϕ is a real-valued convex function on $\mathbb{R}$ and N is a positive integer. Then the function*

$$g(\lambda) = \sum_{i \geq 1} \phi(\lambda_i)$$

is Schur-convex on partitions of N.

Given a partition $\mu = (\mu_1, \mu_2, \ldots, \mu_N)$ with $\mu_i > \mu_j$, the transformation that takes μ to $\rho = (\rho_1, \rho_2, \ldots, \rho_N)$ defined by

$$\begin{aligned} \rho_i &= \mu_i - 1 \\ \rho_j &= \mu_j + 1 \\ \rho_k &= \mu_k, \qquad\qquad k \neq i, j, \end{aligned}$$

is called a *transfer* from i to j. By a result of Muirhead, if $\lambda \prec \mu$, then λ can be derived from μ by successive application of a finite number of transfers ([21], Ch. 5, D.1; [20], (1.16)).

Now consider a partition $\lambda \in \mathbb{P}(m, n)$ where $k_1 = m \leq n = k_2$, which corresponds to a compressed set in $H^2 = K_m \times K_n$. Let $\partial_m \lambda$ and $\partial_n \lambda$ be sets of horizontal and vertical boundary edges of λ, respectively. Then we have

$$|\partial_m \lambda| = \sum_{\lambda_i > 0} \lambda_i (m - \lambda_i) \qquad \textit{and} \qquad |\partial_n \lambda| = \sum_{\lambda'_j > 0} \lambda'_j (n - \lambda'_j)$$

After substituting these in (1) and eliminating constant terms, we see that finding a subset $\lambda \in K_m \times K_n$ minimizing $B(\lambda)$ as defined in (1), is equivalent to *maximizing* the following function $f = f_{m,n}$

$$\begin{aligned} f(\lambda) &= c_1 \sum_{i=1}^{\lambda'_1} \lambda_i^2 + c_2 \sum_{j=1}^{\lambda_1} \lambda_j'^2 \\ &= \gamma_n \sum_{i \geq 1} \lambda_i^2 + \gamma_m \sum_{j \geq 1} \lambda_j'^2 \end{aligned} \tag{4}$$

on $\mathbb{P}(m, n)$ $(m \leq n)$, where

$$\gamma_k = \begin{cases} k^2 & \textit{if } k \textit{ is even}, \\ k^2 - 1 & \textit{if } k \textit{ is odd} \end{cases}$$

in accordance with the definition of the weights c_i in (2).

Note that the function f is not Schur-convex. In other words, transfer operators ([21], Ch. 5, D.1) or equivalently raising/lowering operators (see [20], (1.15-1.16)) which move from a given λ to a smaller one in the linear order while keeping the value of f nondecreasing are insufficient to prove our result[1]. As an example take $m = 4$, $n = 8$, $\lambda = (4, 1, 1, 1, 1, 1, 1)$, $\mu = (4, 2, 1, 1, 1, 1)$. Then $\lambda \prec \mu$ by a single transfer as shown in Figure 3, but $2240 = f(\lambda) > f(\mu) = 2208$.

[1] Transfer arguments can be used to prove Lemma 1 in the special case when $n \geq m^2$.



Figure 3: μ is obtained from λ by single transfer (lowering the indicated cell). Here $m = 4$, $n = 8$, $\lambda \prec \mu$ but $f(\lambda) = 2240$, whereas $f(\mu) = 2208$.

2.1 Extremal sets for the 2–dimensional Hamming graph

Now we are ready to prove the 2–dimensional case which is stated using the terminology of Hamming graphs in the following lemma.

Lemma 1 *Given a 2–dimensional Hamming graph $H^2 = K_m \times K_n$ with $m \leq n$, let λ be any k–vertex subset of $V(H^2)$ and $\overline{\lambda}$ be the set of first k vertices of H^2 in reverse–lexicographic order. Then $f(\overline{\lambda}) \geq f(\lambda)$. That is*

$$\gamma_n \sum_{i \geq 1} \overline{\lambda}_i^2 + \gamma_m \sum_{j \geq 1} \overline{\lambda}_j'^2 \geq \gamma_n \sum_{i \geq 1} \lambda_i^2 + \gamma_m \sum_{j \geq 1} \lambda_j'^2 \tag{5}$$

Proof We only give the proof for n and m both even. The other cases are similar. After rearranging inequality (5), we are required to prove

$$n^2 \sum_{i \geq 1} (\overline{\lambda}_i^2 - \lambda_i^2) \geq m^2 \sum_{j \geq 1} (\lambda_j'^2 - \overline{\lambda}_j'^2) \tag{6}$$

We shall prove the claim by induction on n and m for any given $m \leq n$. The base of the induction is the case where $n = m$, which is already known to be true by Lindsey's result as remarked earlier.

The proof of the inductive step involves different cases based on the number of elements of λ which are in the topmost and bottommost rows of H^2. We call the i^{th} row (from the bottom) of H^2 'empty' with respect to a partition λ if no vertices in this row is a member of λ, and 'nonempty' otherwise. In terms of parts of λ these two conditions mean $\lambda_i = 0$ and $\lambda_i > 0$, respectively. Likewise, the i^{th} row of H^2 is called 'complete' if all vertices in this row belong to the partition (i.e. $\lambda_i = m$), and 'incomplete' ($\lambda_i < m$) otherwise. With this terminology, the proof is divided into the following three cases with respect to λ, covering all possible configurations:

Case (1): empty top row (i.e. $\lambda_n = 0$),
Case (2): complete bottom row (i.e. $\lambda_1 = m$),

Case (3): nonempty top and incomplete bottom rows (i.e. $\lambda_n > 0$ and $\lambda_1 < m$).

We will prove inequality (6) for each one of these cases.

Case (1): *Empty top row*

Assume that the claim is true for $n-1$ where $n-1 \geq m$. Since n is even, $n-1$ is odd, and hence $\gamma_{n-1} = (n-1)^2 - 1 = n^2 - 2n$ according to Definition 1. In this case, the partition λ fits into the 2–dimensional Hamming graph $K_m \times K_{n-1}$ obtained by discarding the top row of H^2, i.e. $\lambda \in \mathbb{P}(m, n-1)$. Thus the induction hypothesis is applicable and we have

$$(n^2 - 2n) \sum_{i=1}^{n-1} (\overline{\lambda}_i^2 - \lambda_i^2) \geq m^2 \sum_{j \geq 1} (\lambda_j'^2 - \overline{\lambda}_j'^2) \tag{7}$$

Combining inequality (6) and (7), it suffices to prove

$$n^2 \sum_{i \geq 1} (\overline{\lambda}_i^2 - \lambda_i^2) \geq (n^2 - 2n) \sum_{i \geq 1} (\overline{\lambda}_i^2 - \lambda_i^2) \tag{8}$$

since $\lambda_n = \overline{\lambda}_n = 0$. After cancellation of terms, this reduces to proving

$$\sum_{i \geq 1} \overline{\lambda}_i^2 \geq \sum_{i \geq 1} \lambda_i^2.$$

Since $|\lambda| = |\overline{\lambda}|$ and the function $\phi(x) = x^2$ is convex on $\mathbb{R}$, this inequality follows from the majorization result of Proposition 1 .

An alternate proof of the last inequality can be reached by the following observation, which further simplifies the proof by eliminating the majorization argument: By combining pairs of rows of λ into a single row if possible (if not, then make one row with size m and another row having the leftover elements) while maintaining the partition property, we eventually convert λ into $\overline{\lambda}$ and do not decrease the function $\sum_{i \geq 1} \lambda_i^2$ at any intermediate step.

Remark
Note that we have not made use of the fact that $\overline{\lambda}$ is in reverse–lexicographic order in the proof of case (1). We have merely used $\lambda \prec \overline{\lambda}$. Therefore when the top row of λ is empty and the ordering ρ is such that $\lambda \prec \rho$, then $f(\lambda) \leq f(\rho)$ by the same proof.

In the proof of case (3) below we give one such ordering.

Case (2): *Complete bottom row*

Consider H^2 and the given configuration λ residing in H^2 with a complete bottommost row. By removing this row from H^2 we obtain a 2–dimensional Hamming graph $K_m \times K_{n-1}$ and a new partition π containing all rows of λ but the bottommost one with $\pi \in \mathbb{P}(m, n-1)$. Let $\overline{\pi}$ be the set of first $|\pi| = |\lambda| - m$ vertices of $K_m \times K_{n-1}$ in reverse–lexicographic order. Note that the first $|\lambda|$ vertices of $K_m \times K_n$ in reverse–lexicographic order are the bottommost row of m elements, followed by $\overline{\pi}$.

Now by the induction hypothesis, we have $f(\overline{\pi}) \geq f(\pi)$, which can be written as

$$(n^2 - 2n) \sum_{i \geq 2} (\overline{\lambda}_i^2 - \lambda_i^2) \geq m^2 \sum_{j \geq 1} (\pi_j'^2 - \overline{\pi}_j'^2) = m^2 \sum_{j \geq 1} (\lambda_j'^2 - \overline{\lambda}_j'^2)$$

In view of the inequality (6) we wish to prove, it suffices to show

$$n^2 \sum_{i \geq 1} (\overline{\lambda}_i^2 - \lambda_i^2) \geq (n^2 - 2n) \sum_{i \geq 2} (\overline{\lambda}_i^2 - \lambda_i^2)$$

which is equivalent to

$$n^2 \sum_{i \geq 1} (\overline{\lambda}_i^2 - \lambda_i^2) \geq (n^2 - 2n) \sum_{i \geq 1} (\overline{\lambda}_i^2 - \lambda_i^2)$$

since $\overline{\lambda}_1^2 = \lambda_1^2 = m^2$. But this is the same inequality as (8) which we already proved in case (1).

Case (3): *Nonempty top and incomplete bottom rows*

Unfortunately, the method used in the previous two cases cannot be applied here, as no row is either completely empty or completely full, and the induction step fails to produce the desired reverse–lexicographic order configuration.

Thus, we use a different approach. Specifically, we convert the given configuration λ into another, say π, where the top row of π is empty and $f(\pi) \geq f(\lambda)$. Evidently, once this is achieved, case (1) above implies that $f(\overline{\lambda}) \geq f(\pi)$ and lemma 1 follows.

We rewrite $f(\lambda)$ of (4) in a form which is more convenient for our treatment. Given a partition $\lambda \in \mathbb{P}(m, n)$, let ω be the partition in $\mathbb{P}(m-1, n-1)$ which is obtained by removing the bottommost row and the leftmost column from the $m \times n$ rectangle. Then $\omega_i = \lambda_{i+1} - 1$ for $1 \leq i \leq n-1$.

Now consider the term $\gamma_n \sum \lambda_i^2$ of $f(\lambda)$ in (4). By taking the first term λ_1^2 out of the summation and putting $\lambda_i = (\lambda_i - 1) + 1$, we have

$$\begin{aligned}
\gamma_n \sum_{i \geq 1} \lambda_i^2 &= \gamma_n \left(\lambda_1^2 + \sum_{i=2}^{\lambda_1'} ((\lambda_i - 1) + 1)^2 \right) \\
&= \gamma_n \left(\lambda_1^2 + \sum_{i \geq 1} \omega_i^2 + 2 \sum_{i \geq 2} \lambda_i - \sum_{i=2}^{\lambda_1'} 1 \right) \\
&= \gamma_n \left(\lambda_1^2 + \sum_{i \geq 1} \omega_i^2 + 2(|\lambda| - \lambda_1) - (\lambda_1' - 1) \right)
\end{aligned}$$

Rewriting the second term $\gamma_m \sum_{j \geq 1} \lambda_j'^2$ of $f(\lambda)$ as above, we now have

$$f(\lambda) = \gamma_n \left(\lambda_1^2 + \sum_{i \geq 1} \omega_i^2 + 2(|\lambda| - \lambda_1) - (\lambda_1' - 1) \right) + \gamma_m \left(\lambda_1'^2 + \sum_{j \geq 1} \omega_j'^2 + 2(|\lambda| - \lambda_1') - (\lambda_1 - 1) \right)$$

Since $f(\omega) = \gamma_n \sum_{i \geq 1} \omega_i^2 + \gamma_m \sum_{j \geq 1} \omega_j'^2$, this is equivalent to

$$f(\lambda) = f(\omega) + \gamma_n \left(\lambda_1^2 + 2(|\lambda| - \lambda_1) - (\lambda_1' - 1) \right) + \gamma_m \left(\lambda_1'^2 + 2(|\lambda| - \lambda_1') - (\lambda_1 - 1) \right) \qquad (9)$$



Figure 4: Obtaining π from λ. Elements in the first row/column are shaded.

Note that, by characterizing $f(\lambda)$ as in equation (9), we have achieved separation of the elements of λ that are in ω from the rest (i.e., those in the first row and the first column). In other words, the contribution of ω to $f(\lambda)$ is now a single additive term $f(\omega)$.

Now the aforementioned configuration π is obtained from λ, as shown in Figure 4, by first moving the $m-\lambda_1$ upper elements in the first column to the end of the first row (thereby $\pi_1 = m$), and next putting the elements corresponding to ω into the reverse–lexicographic order with respect to the $(m-1) \times (n-1)$ rectangle in which it resides.

First we must establish that this resulting configuration is always a partition. This can be done by merely showing that $\pi'_1 \geq \pi'_2$. Evidently, $\pi'_1 = n - m + \lambda_1$. Furthermore, $|w| \leq (\lambda_1 - 1)(n-1)$, and since $|w|$ elements are split into groups of $m-1$ as much as possible due to the nature of reverse–lexicographic order, the following must hold for π'_2.

$$\pi'_2 \leq \left\lceil \frac{(\lambda_1 - 1)(n-1)}{m-1} \right\rceil + 1$$

Thus we are required to prove,

$$n - m + \lambda_1 - 1 \geq \left\lceil \frac{(\lambda_1 - 1)(n-1)}{m-1} \right\rceil$$

Eliminating the *ceiling* from the inequality above, it suffices to prove

$$n - m + \lambda_1 - 1 \geq \frac{(\lambda_1 - 1)(n-1)}{m-1}$$

as the left-hand side is an integer. Multiplying both sides by $(m-1)$ and rearranging terms yield $m(n-m) \geq \lambda_1(n-m)$ which obviously holds as $m \geq \lambda_1$ and $n \geq m$. Thus, π will always be a partition.

As in the construction of ω, let μ be the partition in $\mathbb{P}(m-1, n-1)$ obtained by removing the bottommost row and the leftmost column from the $m \times n$ rectangle where π resides. Now we would like to show $f(\pi) \geq f(\lambda)$, which is equivalent to showing

$$f(\mu) + n^2\left(\pi_1^2 + 2(|\pi| - \pi_1) - (\pi'_1 - 1)\right) + m^2\left(\pi_1'^2 + 2(|\pi| - \pi'_1) - (\pi_1 - 1)\right) \geq \tag{10}$$

$$f(\omega) + n^2\Big(\lambda_1^2 + 2(|\lambda| - \lambda_1) - (\lambda_1' - 1)\Big) + m^2\Big(\lambda_1'^2 + 2(|\lambda| - \lambda_1') - (\lambda_1 - 1)\Big)$$

in view of (9).

An important point to note is that, due to the term $f(\omega)$ in equation (9), the partition ω (and μ) can be considered as being in the $m \times n$ rectangle in terms of its contribution to $f(\lambda)$ besides the $(m-1) \times (n-1)$ one in which it is originally defined. This is because the function f is determined in equation (4) with respect to the $m \times n$ rectangle. Therefore the top row of $\omega \in \mathbb{P}(m,n)$ is empty. However μ is not in reverse–lexicographic order with respect to the $m \times n$ rectangle, as $\mu_m' = 0$. Fortunately, $\omega \prec \mu$, and therefore we have $f(\omega) \leq f(\mu)$ by the remark made at the end of case (1). Thus to establish inequality (10), it suffices to prove the following.

$$n^2\Big(\pi_1^2 + 2(|\pi| - \pi_1) - (\pi_1' - 1)\Big) + m^2\Big(\pi_1'^2 + 2(|\pi| - \pi_1') - (\pi_1 - 1)\Big) \geq$$

$$n^2\Big(\lambda_1^2 + 2(|\lambda| - \lambda_1) - (\lambda_1' - 1)\Big) + m^2\Big(\lambda_1'^2 + 2(|\lambda| - \lambda_1') - (\lambda_1 - 1)\Big)$$

Using $|\pi| = |\lambda|$ and eliminating the constant terms, the inequality above follows iff

$$n^2\pi_1^2 - n^2(2\pi_1 + \pi_1') + m^2\pi_1'^2 - m^2(2\pi_1' + \pi_1) \geq n^2\lambda_1^2 - n^2(2\lambda_1 + \lambda_1') + m^2\lambda_1'^2 - m^2(2\lambda_1' + \lambda_1)$$

Since $\lambda_1 + \lambda_1' = \pi_1 + \pi_1'$, $\pi_1 = m$ and $\lambda_1' = n$, this holds iff

$$n^2(m + \lambda_1 - 1) \geq m^2(n + \pi_1' - 1)$$

Using the fact that $\pi_1' = n - m + \lambda_1$ and factorizing, we have

$$(n-m)(nm - m^2) + (n^2 - m^2)(\lambda_1 - 1) \geq 0$$

which obviously holds for $n \geq m$. □

The lemma is proved. Now we are ready to prove the main theorem for higher dimensions.

3 The higher–dimensional case

In this section, we prove Theorem 1 for an arbitrary number of dimensions d. The main idea of the proof is based on that of [16], hence our notation is similar to the notation therein.

Proof The proof is by induction on d with $d = 2$, already proved in Lemma 1, being the base case. We assume $k_1 \leq k_2 \leq \cdots \leq k_d$ for $H^d = K_{k_1} \times K_{k_2} \times \cdots \times K_{k_d}$ and vertices are labelled by d-tuples $\langle l_1, l_2, \cdots, l_d \rangle$ where $0 \leq l_i \leq k_i - 1$.

The idea is to transform a given arbitrary configuration into one in reverse–lexicographic order so as not to increase the normalized boundary. To aid the readability of the proof, Figure 5 provides a 3–dimensional Hamming graph $H^3 = K_4 \times K_5 \times K_{10}$ which illustrates the transformation process.

Given an arbitrary configuration X in H^d, we permute the k_d $(d-1)$–dimensional Hamming subgraphs along dimension d such that successive subgraphs have fewer elements of X. Now we apply the induction hypothesis to each of these subgraphs. Phase (i) in Figure 5 illustrates a

configuration obtained after this step. Note that applying this procedure cannot increase $B(X)$ since $|\partial_d X|$ cannot increase and by the induction hypothesis

$$\frac{|\partial_1 X_i|}{c_1} + \cdots + \frac{|\partial_{d-1} X_i|}{c_{d-1}}$$

is smallest for each subgraph i, where X_i is the set of elements of X that are in subgraph i. Not surprisingly, this means that candidate extremal sets in higher dimensions are among *higher–dimensional partitions* (see [2], Ch. 11), which are contained in the d-dimensional parallelepiped $k_1 \times k_2 \times \cdots \times k_d$. Now we repeat the same steps for subgraphs along dimension $d-1$ as well. This step is illustrated by phase (ii) in Figure 5.

Consider the $(d-2)$–dimensional Hamming subgraphs of H^d when dimensions d and $d-1$ are fixed. We call each such subgraph 'complete' iff its vertices are completely contained in X, 'incomplete' iff there exists some (but not all) contained in X and 'empty' iff none is in X. We shall show that if there are more than one incomplete subgraphs then these can be combined without increasing $B(X)$. The result of this step is shown by phase (iii) in Figure 5. To this end we give some definitions and develop proper notation.



Figure 5: Conversion into the reverse–lexicographic order in three dimensions.

First, suppose $P^*_{p,q}$ and $P^*_{r,s}$ are sets of vertices of two such incomplete $(d-2)$–dimensional Hamming subgraphs where p,r and q,s are coordinates of dimension d and $d-1$ respectively with $0 \le p, r \le k_d - 1$ and $0 \le q, s \le k_{d-1} - 1$. Without loss of generality, assume

$$\frac{p}{c_d} + \frac{q}{c_{d-1}} \ge \frac{r}{c_d} + \frac{s}{c_{d-1}} \tag{11}$$

Next, let $P_{p,q} = P^*_{p,q} \cap X$, $P_{r,s} = P^*_{r,s} \cap X$ and $Y = X \setminus (P_{p,q} \cup P_{r,s})$. In our example, there are exactly two such subgraphs with $p = 0$, $q = 3$ and $r = 2$, $s = 2$, i.e. $P_{0,3}$ and $P_{2,2}$, which satisfies the assumption given by the inequality above.

Now given two disjoint subsets S and T of $V(H^d)$, let

$$B(S,T) = \frac{|\partial_1(S,T)|}{c_1} + \frac{|\partial_2(S,T)|}{c_2} + \cdots + \frac{|\partial_d(S,T)|}{c_d} \tag{12}$$

where c_i is as defined before and $\partial_i(S,T)$ is the set of edges in dimension i having one end in S and the other in T. Note that, in this notation $B(X) = B(X, V(H^d) \setminus X)$.

Note that the following holds.

$$B(X) = B(Y) + B(P_{p,q}) + B(P_{r,s}) - 2(B(Y, P_{p,q}) + B(Y, P_{r,s})) - 2B(P_{p,q}, P_{r,s}) \tag{13}$$

We claim that if as many elements in $P_{r,s}$ as possible are moved to $P^*_{p,q}$ preserving the reverse–lexicographic order then $B(X)$ does not increase. To this end, consider the terms in equation (13). First we remark that, by virtue of the reverse–lexicographic order, we must have $p \neq r$ and $q \neq s$, and therefore $B(P_{p,q}, P_{r,s}) = 0$ in equation (13). Furthermore, because of inequality (11), $B(Y, P_{p,q}) + B(Y, P_{r,s})$ cannot decrease by this move, and $B(Y)$ is constant. Finally, we claim that $B(P_{p,q}) + B(P_{r,s})$ does not increase.

To prove this, note that any vertex $v \in (P_{p,q} \cup P_{r,s})$ is adjacent to $k_d - 1$ and $k_{d-1} - 1$ vertices in dimensions d and $d-1$, respectively. Thus, moving vertices from $P_{r,s}$ to $P^*_{p,q}$ does not change the boundary along dimensions d and $d-1$. Therefore, it suffices to prove

$$B'(P_{p,q}) + B'(P_{r,s}) \geq B'(P'_{p,q}) + B'(P'_{r,s}) \tag{14}$$

where

$$B'(X) = \frac{|\partial_1 X|}{c_1} + \cdots + \frac{|\partial_{d-2} X|}{c_{d-2}}$$

and $P'_{p,q}$ and $P'_{r,s}$ are the new subsets corresponding to $P_{p,q}$ and $P_{r,s}$ respectively, after elements are moved from $P_{r,s}$ to $P^*_{p,q}$.

To prove inequality (14), first suppose that all of $P_{r,s}$ fits in the complement of $P_{p,q}$ with respect to $P^*_{p,q}$. Thus we can place elements of $P_{r,s}$ into $P^*_{p,q} \setminus P_{p,q}$ in a set structurally identical to the one given by reverse–lexicographic order, i.e. starting with vertex $\langle k_1 - 1, k_2 - 1, \cdots, k_{d-2} - 1, q, p\rangle$ of H^d and expanding in the direction of the smallest factor of the Hamming graph. That is

$$\langle k_1 - 1, k_2 - 1, \cdots, k_{d-3} - 1, k_{d-2} - 1, q, p\rangle \to \langle k_1 - 1, k_2 - 1, \cdots, k_{d-3} - 1, k_{d-2} - 2, q, p\rangle \to \ldots$$

$$\ldots \to \langle k_1 - 1, k_2 - 1, \cdots, k_{d-3} - 2, k_{d-2} - 1, q, p\rangle \to \langle k_1 - 1, k_2 - 1, \cdots, k_{d-3} - 2, k_{d-2} - 2, q, p\rangle \to \ldots$$

and so on. This is shown in Figure 6.

In this case, we have $B'(P'_{r,s}) = 0$ since $P'_{r,s} = \phi$ and $B'(P'_{p,q})$ can be written as $B'(P'_{p,q}) = B'(P_{p,q}) + B'(P_{r,s}) - 2B'(P_{p,q}, P_{r,s})$. Substituting these values in inequality (14), it suffices to prove

$$B'(P_{p,q}) + B'(P_{r,s}) \geq B'(P_{p,q}) + B'(P_{r,s}) - 2B'(P_{p,q}, P_{r,s})$$

which obviously holds since $B'(P_{p,q}, P_{r,s}) \geq 0$. We remark that $P'_{p,q}$ is not in reverse–lexicographic order at this point since it consists of two subsets each of which is structurally in reverse–lexicographic order. Nevertheless, by an easy application of the induction hypothesis, we can convert it to the reverse–lexicographic order without increasing $B'(P'_{p,q})$.

Now assume that not all elements of $P_{r,s}$ fit into $P^*_{p,q}$. Take first $|P^*_{p,q} \setminus P_{p,q}|$ vertices in reverse–lexicographic order in $P^*_{r,s}$. These vertices are in $P_{r,s}$. Call this set of vertices Y_2 and set $Y_1 = P_{r,s} \setminus Y_2$. After moving all vertices in Y_2 to $P^*_{p,q}$, we put Y_1 in reverse–lexicographic order $\overline{Y}_1$ within $P^*_{r,s}$. This is shown in Figure 7.

Then, inequality (14) reduces to proving

$$B'(P_{p,q}) + B'(P_{r,s}) \geq B'(Y_1)$$



Figure 6: Combining two incomplete subgraphs where the elements can fit into one. (i) subgraphs before, (ii) after.



Figure 7: Combining two incomplete subgraphs where the elements cannot fit into one. (i) Subgraphs before, (ii) after.

as $B'(Y_1) \geq B'(\overline{Y_1})$ holds by the induction hypothesis. Now note that $B'(P_{r,s}) = B'(Y_1)+B'(Y_2)-2B'(Y_1,Y_2)$ and $B'(Y_2) = B'(P_{p,q})$ since Y_2 and $P_{p,q}$ are complementary in $P^*_{p,q}$. Thus the above inequality is equivalent to

$$B'(Y_2) \geq B'(Y_1, Y_2)$$

which obviously holds. Thus $B(P_{p,q}) + B(P_{r,s})$ does not increase as claimed. By applying this process to all $(d-2)$–dimensional incomplete subgraphs, we can assume that X has only one incomplete $(d-2)$–dimensional Hamming subgraph.

Finally we treat the $(d-2)$–dimensional Hamming subgraphs as single vertices and use the 2–dimensional case to minimize $B(X)$ by putting them in reverse–lexicographic order with the only incomplete one highest in the order, as shown by phase (iv) in Figure 5. This completes the proof of Theorem 1. □

4 Conclusions

We proved that the set of first k vertices of the Hamming graph $H^d = K_{k_1} \times K_{k_2} \times \cdots \times K_{k_d}$ $(k_1 \leq k_2 \leq \cdots \leq k_d)$, in reverse–lexicographic order constitutes an extremal set minimizing the dimension–normalized edge–boundary over all k–vertex subsets of the graph. The boundary edges $\partial_i X$ along the i-th dimension of $X \subset V(H^d)$ are normalized by a weight

$$c_i = \begin{cases} k_i^2 & if\ k_i\ is\ even, \\ k_i^2 - 1 & if\ k_i\ is\ odd, \end{cases}$$

which naturally arises in the isoperimetric number problem for d-dimensional arrays. The weighted boundary to be minimized is then

$$B(X) = \frac{|\partial_1 X|}{c_1} + \frac{|\partial_2 X|}{c_2} + \cdots + \frac{|\partial_d X|}{c_d}$$

over $X \subset V(H^d)$. Interestingly, when all factors of H^d have equal size, the lexicographic and reverse–lexicographic orders both result in structurally symmetric subsets and hence are equivalent with respect to extremal sets minimizing the boundary (dimension–normalized or otherwise). Thus our result is identical to Lindsey's for $k_1 = k_2 = \cdots = k_d$.

We formulated the problem for the 2-dimensional case as the maximization of the function f defined on partitions $\lambda \in \mathbb{P}(m,n)$, $(m \leq n)$, by

$$f(\lambda) = \gamma_n \sum_{i=1}^{n} \lambda_i^2 + \gamma_m \sum_{j=1}^{m} \lambda_j'^2$$

and proved that f is maximized for $N \leq nm$, by the reverse–lexicographically smallest partition of N in $\mathbb{P}(m,n)$, where

$$\gamma_n = \begin{cases} n^2 & if\ n\ is\ even, \\ n^2 - 1 & if\ n\ is\ odd \end{cases}$$

This result for $d = 2$ forms the base step of the higher dimensional case.

In light of the remark following the proof of case (1) and the fact that a similar one exists for case (2) in the two–dimensional proof, one may consider the following generalization of Lemma 1: In two dimensions $m \times n$, $m \leq n$, the normalized–boundary of the set of first k vertices in r–limited reverse–lexicographic order (where each row has at most $r \leq m$ elements) is smaller than that of any configuration λ with $\lambda_1 \leq r$. However, this is incorrect. As counterexample, take $\lambda = (2,2,2,2,2,2,2,2)$ in 6×8 rectangle where the set of first $|\lambda| = 16$ vertices in 4–limited reverse–lexicographic order is given by $\overline{\lambda} = (4,4,4,4)$, but $B(\overline{\lambda}) \not\leq B(\lambda)$. However, a similar version of this statement could be true when the normalization for boundary along each dimension is a quadratic function not of dimension itself but the maximum elements in rows or columns in a given configuration, such as r above.

In fact it would be interesting to see a characterization of functions θ_n for which

$$f(\lambda) = \frac{\theta_n}{\theta_m} \sum_{i=1}^{n} \lambda_i^2 + \sum_{j=1}^{m} \lambda_j'^2$$

is maximized by the reverse–lexicographically smallest partition of N in $\mathbb{P}(m,n)$.

Acknowledgement

The authors would like to thank the anonymous referee whose careful review contributed greatly to the robustness of the proofs given in this revised version.

References

[1] R. Ahlswede and S.L. Bezrukov. "Edge-isoperimetric theorems for integer point arrays," *Appl. Math. Letters,* **8**(2), pp. 75–80, 1995.

[2] G.E. Andrews. *The Theory of Partitions,* Addison–Wesley Publishing Company, Reading, MA, 1976.

[3] M. C. Azizoğlu and Ö. Eğecioğlu. "Isoperimetric number of the Cartesian product of graphs and paths," *Congressus Numerantium,* **131**, pp. 135–143, 1998.

[4] M. C. Azizoğlu and Ö. Eğecioğlu. "The isoperimetric number of d–dimensional k–ary arrays," *Int. J. of Found. of Computer Sci.,* **10**(3), pp. 289–300, 1999.

[5] M. C. Azizoğlu and Ö. Eğecioğlu. "The isoperimetric number of products of paths and cycles," Proceedings of the 9^{th} Quadrennial Int. Conf. on Graph Theory, Combinatorics, Algorithms and Applications (GTCC2000), Western Michigan University, June 2000.

[6] M. C. Azizoğlu and Ö. Eğecioğlu. "The bisection width and the isoperimetric number of arrays," Technical Report TRCS00-12, Department of Computer Science, University of California Santa Barbara, June 2000.

[7] S.L. Bezrukov. "Variational principle in discrete extremal problems," *Technical Report,* TR–RI 94–152, Univ. of Paderborn, Paderborn, Germany, 1994.

[8] S.L. Bezrukov. "Edge isoperimetric problems on graphs," *Graph Theory and Combinatorial Biology,* Bolyai Soc. Math. Stud. 7, L. Lovasz, A. Gyarfas, G.O.H. Katona, A. Recski, L. Szekely, Eds., Budapest, 1999, pp. 157–197.

[9] B. Bollobás. *Combinatorics* Cambridge University Press, 1986.

[10] B. Bollobás and I. Leader. "An isoperimetric inequality on the discrete torus," *SIAM J. Disc. Math.,* **3**(1), pp. 32–37, 1990.

[11] B. Bollobás and I. Leader. "Edge-isoperimetric inequalities in the grid," *Combinatorica,* **11** (4), pp. 299–314, 1991.

[12] F.R.K. Chung. *Spectral Graph Theory,* AMS Regional Conference Series in Mathematics, **92**, 1997.

[13] F.R.K. Chung and P. Tetali. "Isoperimetric inequalities for Cartesian products of graphs," *Combinatorics, Probability & Computing,* **7**(2), pp. 141–148, 1998.

[14] L.H. Harper. "Optimal assignment of numbers to vertices," *J. Soc. Ind. Appl. Math.,* **12**(1), pp. 131–135, 1964.

[15] L.H. Harper. "On an isoperimetric problem for Hamming graphs," *Discrete Applied Mathematics,* **95**(1–3), pp. 285–309, 1999.

[16] D.J. Kleitman, M.M. Krieger and B.L. Rothschild. "Configurations maximizing the number of pairs of Hamming–adjacent lattice points," *Studies in Applied Mathematics,* **1**(2), pp. 115–119, 1971.

[17] I. Leader, *Personal communication.*

[18] F.T. Leighton. *Introduction to Parallel Algorithms and Architectures: Arrays · Trees · Hypercubes.* Morgan Kaufmann Publishers, San Mateo, California, 1992.

[19] J.H. Lindsey II. "Assignment of numbers to vertices," *Amer. Math. Monthly,* **7**, pp. 508–516, 1964.

[20] I.G. Macdonald. *Symmetric Functions and Hall Polynomials*, Second edition, Clarendon Press, Oxford, 1995.

[21] Marshall A. W., and I. Olkin. *Inequalities: Theory of Majorization and Its Applications*, Academic Press Inc., San Diego, 1979.

[22] B. Mohar. "Isoperimetric numbers of graphs," *J. of Comb. Theory,* Series B **47**, pp. 274–291, 1989.

[23] K. Nakano. "Linear layouts of generalized hypercubes," *Graph-Theoretic Concepts in Computer Science,* (Jan van Leeuwen Ed.), Springer-Verlag LNCS, **790**, pp. 364–375, 1993.

[24] O. Riordan. "An ordering on the even discrete torus," *SIAM J. Disc. Math.,* **11**(1), pp. 110–127, 1998.

[25] D. Wang and P. Wang. "Discrete isoperimetric problems," *SIAM J. Appl. Math.,* **32**(4), pp. 860–870, 1977.